

Index to Volume 74

Author Index

- Abu-Tarboush, H. M., S. A. B. Ahmed, and H. A. Al Kahtani. Note: Some nutritional and functional properties of karkade (*Hibiscus sabdariffa*) seed products, 352
- Adalsteins, A. E. See W. H. Vensel, 108
- Aguerre, R. J. See M. P. Tolaba, 842
- Ahmed, S. A. B. See H. M. Abu-Tarboush, 352
- Aime, D. B. See H. Liu, 159
- Al Kahtani, H. A. See H. M. Abu-Tarboush, 352
- Anderson, B. J. See C. M. McDonough, 542
- Andrews, L. C. See C. S. Gaines, 700
- Apintanapong, M. See A. Nookhorm, 12
- Arendt, E. K. See K. Wehrle, 739
- Armfield, S. D. See H. Liu, 159
- Autio, K. See J. Räsänen, 56, 806
- Au-Yeung, L. See H. Corke, 356
- Baik, B.-K. See T. Otto, 141, 445
- Bailey, T. B. See S. K. Singh, 40
- Barreiro, N. L. See V. Singh, 672
- Barrett, J. E., C. F. Klopfenstein, and H. W. Leipold. Detoxification of rapeseed meal by extrusion with an added basic salt, 168
- Barton, F. E. II See E. T. Champagne, 566
- See W. R. Windham, 626
- Batey, I. L., B. M. Curtin, and S. A. Moore. Optimization of rapid-visco analyser test conditions for predicting Asian noodle quality, 497
- See M. Zeng, 63
- Bean, S. R., and G. L. Lookhart. Separation of wheat proteins by two-dimensional reversed-phase high-performance liquid chromatography plus free zone capillary electrophoresis, 758
- Bechtel, D. B., and J. D. Wilson. Ultrastructure of developing hard and soft red winter wheats after air- and freeze-drying and its relationship to endosperm texture, 235
- Beer, M. U., P. J. Wood, and J. Weisz. Molecular weight distribution and (1→3)(1→4)- β -D-glucan content of consecutive extracts of various oat and barley cultivars, 476
- , —, —, and N. Fillion. Effect of cooking and storage on the amount and molecular weight of (1→3)(1→4)- β -D-glucan extracted from oat products by an in vitro digestion system, 705
- Beléa, A. P. See M. A. Larrea, 98
- BeMiller, J. N. See K. C. Huber, 537
- See M. R. Jacobson, 511
- See M. Obanni, 431
- Bergman, E.-L. See A. Rutgersson, 727
- Bett, K. L. See E. T. Champagne, 566
- Bhattacharya, M., J. Jafari-Shabestari, C. O. Qualset, and H. Corke. Diversity of starch pasting properties in Iranian hexaploid wheat landraces, 417
- Bhatty, R. S. Milling of regular and waxy starch hull-less barleys for the production of bran and flour, 693
- , and B. G. Rossnagel. Communication to the editor: Zero amylose lines of hull-less barley, 190
- See T. Vasanathan, 25
- Bietz, J. A. See F. R. Huebner, 123
- Blakeney, A. B. See B. G. Osborne, 467
- Blanshard, J. M. V. See J. Räsänen, 806
- Borras, F. S. See J. L. Robutti, 75
- Borri, V. See G. Giovanelli, 710
- Botham, R. L., P. Cairns, R. M. Faulks, G. Livesey, V. J. Morris, T. R. Noel, and S. G. Ring. Physicochemical characterization of barley carbohydrates resistant to digestion in a human ileostomate, 29
- Brumm, T. J. See M. R. Campbell, 300
- Bryant, C. M., and B. R. Hamaker. Effect of lime on gelatinization of corn flour and starch, 171
- Buffo, R. A., C. L. Weller, and A. Gennadios. Note: Films from laboratory-extracted sorghum kafirin, 473
- Bullerman, L. B. See S. K. Katta, 858
- Buriak, P. See J. F. Lopes-Filho, 633
- See V. Singh, 672
- Bushuk, W., R. L. Hay, N. G. Larsen, R. G. Sara, L. D. Simmons, and K. H. Sutton. Effect of mechanical dough development on the extractability of wheat storage proteins from bread dough, 389
- Cagampang, A. E. See S. K. Katta, 858
- Cairns, P. See R. L. Botham, 29
- Campbell, M. R., T. J. Brumm, and D. V. Glover. Whole grain amylose analysis in maize using near-infrared transmittance spectroscopy, 300
- Campos, D. T., J. F. Steffe, and P. K. W. Ng. Rheological behavior of undeveloped and developed wheat dough, 489
- Cañigueral, S. See M. Estiarte, 495
- Casals, I. See M. Estiarte, 495
- Çelik, S. See D. Dadayli, 375
- Cerletti, P. See N. Guerrieri, 846
- Champagne, E. T., K. L. Bett, B. T. Vinyard, B. D. Webb, A. M. McClung, F. E. Barton, II, B. G. Lyon, K. Moldenhauer, S. Linscombe, and D. Kohlwey. Effects of drying conditions, final moisture content, and degree of milling on rice flavor, 566
- See H. S. Guraya, 561
- See W. R. Windham, 626
- Chandrasekar, A. See R. Kavitha, 22
- Chang, P. See T. Vasanathan, 25
- Charvat, B. See U. Pechanek, 800
- Chen, H., B. P. Marks, and T. J. Siebenmorgen. Quantifying surface lipid content of milled rice via visible/near-infrared spectroscopy, 826
- , and T. J. Siebenmorgen. Effect of rice kernel thickness on degree of milling and associated optical measurements, 821
- Chen, L.-N. See S. Lu, 34
- Chen, X. See H. Corke, 356
- Cherian, G., and P. Chinachoti. Action of oxidants on water sorption, ^2H nuclear magnetic resonance mobility, and glass transition behavior of gluten, 312
- Chibbar, R. N. See T. Demeke, 442
- Chinachoti, P. See G. Cherian, 312
- Chiu, M.-C. M. See W. H. Yokoyama, 293
- Chiu, M. M. See B. E. Knuckles, 599
- Chung, O. K. See J. L. Hazelton, 400
- See F. R. Huebner, 123
- See H. Park, 202, 207
- Clarke, J. M. See J. E. Dexter, 519
- Clear, R. M. See J. E. Dexter, 519
- Cohen, C. See J. Fu, 304
- Collado, L. S., and H. Corke. Properties of starch noodles as affected by sweetpotato genotype, 182
- , L. B. Mabesa, and —. Genetic variation in color of sweetpotato flour related to its use in wheat-based composite flour products, 681
- Corke, H., L. Au-Yeung, and X. Chen. Note: An automated system for the continuous measurement of time-dependent changes in noodle color, 356
- See M. Bhattacharya, 417
- See L. S. Collado, 182, 681
- See W. Li, 79
- Cortez-Rocha, M. O. See R. I. Sánchez-Mariñez, 481
- Craig, J. C., Jr. See L. C. Dickey, 676
- See N. Parris, 766
- Crosbie, G. B. See A. S. Ross, 814
- Cuppert, S. L. See L. A. Kunte, 115
- Curtin, B. M. See I. L. Batey, 497
- Czuchajowska, Z. See P.-Y. Lin, 639
- See T. Otto, 141, 445
- Dadayli, D., M. M. Sünnetçioğlu, H. Köksel, and S. Çelik. Detection of irradiated wheat using the electron paramagnetic resonance spin probe technique, 375
- Dahl, W. J. See A. M. Stephen, 379
- Daigle, K. See F. F. Shih, 437
- Dallmer, M. F. See L. C. Dickey, 676
- Dejmeck, P. See K. M. Lindborg, 49
- Demeke, T., P. Hucl, R. B. Nair, T. Nakamura, and R. N. Chibbar. Note: Evaluation of Canadian and other wheats for waxy proteins, 442
- Dexter, J. E., B. A. Marchylo, R. M. Clear, and J. M. Clarke. Effect of Fusarium head blight on semolina milling and pasta-making quality of durum wheat, 519
- Dhaliwal, Y. S. See D. W. Hatcher, 502
- Dickey, L. C., M. F. Dallmer, E. R. Radewonuk, N. Parris, M. Kurantz, and J. C. Craig, Jr. Hydrocyclone separation of dry-milled corn, 676
- See N. Parris, 766
- Dintzis, F. R. See J. A. Klavons, 832
- See M. M. Millard, 687
- Doehlert, D. C., and W. R. Moore. Composition of oat bran and flour prepared by three different mechanisms of dry milling, 403

- See D. Zhang, 722
- Dornbrink-Kurtzman, M. A., and C. A. Knutson. A study of maize endosperm hardness in relation to amylose content and susceptibility to damage, 776
- Doner, L. W., and K. B. Hicks. Isolation of hemicellulose from corn fiber by alkaline hydrogen peroxide extraction, 176
- Dowd, M. K. Recovery of starch and protein from wet-milled corn fiber, 589
- Dowell, F. E. Effect of NaOH on visible wavelength spectra of single wheat kernels and color classification efficiency, 617
- Du, L. See S. K. Singh, 254
- Duvick, S. A. See K. M. Krieger, 553
- See K.-Y. Ng, 288, 837
- Eastman, J. J. See J. L. Hazelton, 400
- Eckhoff, S. R. See J. F. Lopes-Filho, 633
- See K. D. Rausch, 274
- See S. K. Mehra, 484
- See S. K. Singh, 254
- See V. Singh, 249, 462, 672
- Eggum, B. O., and B. O. Juliano. Properties and protein quality in growing rats of a low-glutenin content rice mutant, 200
- Eliasson, A.-C. See K. M. Lindborg, 49
- Ellis, R. P. See J. S. Swanston, 456
- Endo, S. See K. Hayakawa, 576
- Englyst, H. N. See A. M. Stephen, 379
- Estiarte, M., J. Peñuelas, S. Cañigueral, and I. Casals. Note: A reverse-phase HPLC method for tricin separation from wheat leaves, 495
- Eyherabide, G. H. See J. L. Robutti, 75
- Eynard, L. See N. Guerrieri, 846
- Fahloul, D. See M. G. Scanlon, 612
- Faulks, R. M. See R. L. Botham, 29
- Fearn, T. See B. G. Osborne, 467
- Fillion, N. See M. U. Beer, 705
- Finney, P. L. See C. S. Gaines, 700
- Fisher, D. K., and D. B. Thompson. Retrogradation of maize starch after thermal treatment within and above the gelatinization temperature range, 344
- Fox, S. R. See S. K. Singh, 40
- See S. Wu, 264
- Fu, J., S. J. Mulvaney, and C. Cohen. Effect of added fat on the rheological properties of wheat flour doughs, 304
- Gaines, C. S., P. L. Finney, and L. C. Andrews. Influence of kernel size and shriveling on soft wheat milling and baking quality, 700
- Garber, B. W., F. Hsieh, and H. E. Huff. Influence of particle size on the twin-screw extrusion of corn meal, 656
- Gelroth, J. A. See G. S. Ranhotra, 297, 361
- Gennadios, A. See R. A. Buffo, 473
- See M. Kojima, 526
- See L. A. Kunte, 115
- Giovanelli, G., C. Peri, and V. Borri. Effects of baking temperature on crumb-staling kinetics, 710
- Glover, D. V. See M. R. Campbell, 300
- Grau, H. See K. Wehrle, 739
- Gröger, S. See U. Pechanek, 800
- Grosch, W., and P. Schieberle. Flavor of cereal products—A review, 91
- Grossmann, M. V. E. See M. A. Larrea, 98
- Guerrieri, N., L. Eynard, V. Lavelli, and P. Cerletti. Interactions of protein and starch studied through amyloglucosidase action, 846
- Guraya, H. S., R. S. Kadan, and E. T. Champagne. Effect of rice starch-lipid complexes on in vitro digestibility, complexing index, and viscosity, 561
- Haginoya, R., K. Sakai, T. Komatsu, S. Nagao, K. Yokoyama, O. Takeuchi, R. Matsukawa, and I. Karube. Determination of damaged starch and diastatic activity in wheat flour using a flow injection analysis biosensor method, 745
- Haken, A. E. See V. Singh, 249
- Hamada, J. S. Characterization of protein fractions of rice bran to devise effective methods of protein solubilization, 662
- Hamaker, B. R. See C. M. Bryant, 171
- Hammond, E. G. See M. Zhang, 594
- Hanna, M. A. See M. Kojima, 526
- See A. N. R. Kollengode, 396
- See L. A. Kunte, 115
- Hatcher, D. W., and J. E. Kruger. Simple phenolic acids in flours prepared from Canadian wheat: Relationship to ash content, color, and polyphenol oxidase activity, 337
- , —, and Y. S. Dhaliwal. Effect of roller mill extraction rate on the chapatti-making quality of Canadian flours, 502
- Hay, R. L. See W. Bushuk, 389
- Hayakawa, K., K. Tanaka, T. Nakamura, S. Endo, and T. Hoshino. Quality characteristics of waxy hexaploid wheat (*Triticum aestivum* L.): Properties of starch gelatinization and retrogradation, 576
- Hayashi, M. See M. Seguchi, 129, 384
- Hazelton, J. L., O. K. Chung, J. J. Eastman, C. E. Lang, P. J. McCluskey, R. A. Miller, M. A. Shipman, and C. E. Walker. Note: Regression equation for predicting absorption for 2-g direct drive mixograph, 400
- Hicks, K. B. See L. W. Doner, 176
- Hix, D. K., C. F. Klopfenstein, and C. E. Walker. Physical and chemical attributes and consumer acceptance of sugar-snap cookies containing naturally occurring antioxidants, 281
- Hizukuri, S. See C. P. Villareal, 163
- Holley, R. A. See H. Liu, 159
- Horsley, R. D. See C. J. Lee, 571
- Hoseney, R. C. See R. A. Miller, 330, 614, 669
- Hoshino, T. See K. Hayakawa, 576
- Hsieh, F. See B. W. Garber, 656
- Huang, D. Y., and K. Khan. Characterization and quantification of native glutenin aggregates by multistacking sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) procedures, 229
- and —. Quantitative determination of high molecular weight glutenin subunits of hard red spring wheat by SDS-PAGE. I. Quantitative effects of total amounts on breadmaking quality characteristics, 781
- and —. Quantitative determination of high molecular weight glutenin subunits of hard red spring wheat by SDS-PAGE. II. Quantitative effects of individual subunits on breadmaking quality characteristics, 786
- Huber, K. C., and J. N. BeMiller. Visualization of channels and cavities of corn and sorghum starch granules, 537
- Hucl, P. See T. Demeke, 442
- Hudson, C. A. See W. H. Yokoyama, 293
- Huebner, F. R., T. C. Nelsen, O. K. Chung, and J. A. Bietz. Protein distributions among hard red winter wheat varieties as related to environment and baking quality, 123
- Huff, H. E. See B. W. Garber, 656
- Hussain, A., and O. M. Lukow. Influence of gliadin-rich subfractions of Glenlea wheat on the mixing characteristics of wheat flour, 791
- , —, B. M. Watts, and R. I. H. McKenzie. Rheological properties of full-formula doughs derived from near-isogenic 1BL/1RS translocation lines, 242
- Ingledeu, W. M. See S. Wang, 621
- Irving, D. W., and I. A. Jideani. Microstructure and composition of *Digitaria exilis* Stapf (acha): A potential crop, 224
- Jackson, L. S. See S. K. Katta, 858
- Jacobson, M. R., M. Obanni, and J. N. BeMiller. Retrogradation of starches from different botanical sources, 511
- Jafari-Shabestari, J. See M. Bhattacharya, 417
- Jideani, I. A. See D. W. Irving, 224
- Jo, J. S. See S. S. Kim, 212
- Johns, D. M. See A. M. Stephen, 379
- Johnson, L. A. See S. K. Singh, 40
- See S. Wu, 258, 264, 268
- Jones, B. L. See N. Zhang, 119
- Juliano, B. O. See B. O. Eggum, 200
- See C. P. Villareal, 163
- Kachman, S. D. See W. J. Park, 7
- Kadan, R. S. See H. S. Guraya, 561
- Kaid, N., L. Rakotozafy, J. Potus, and J. Nicolas. Studies on the glutathione-dehydroascorbate oxidoreductase (EC 1.8.5.1) from wheat flour, 605
- Kanenaga, K. See M. Seguchi, 542
- Karger, A. See U. Pechanek, 800
- Karube, I. See R. Haginoya, 745
- Kasarda, D. D. See P. Köhler, 154
- See W. H. Vensel, 108
- Katta, S. K., A. E. Cagampang, L. S. Jackson, and L. B. Bullerman. Distribution of *Fusarium* molds and fumonisins in dry-milled corn fractions, 858
- Kaukovirta-Norja, A., P. Reinikainen, J. Oikku, and S. Laakso. Starch lipids of barley and malt, 733
- Kavitha, R., and A. Chandrashekar. Content and composition of nonstarchy polysaccharides in endosperms of sorghum varying in hardness during four developmental stages, 22
- Keck-Gassenmeier, B. See P. Köhler, 154
- Khan, K. See D. Y. Huang, 229, 781, 786
- Kidwell, K. K. See C. F. Morris, 16

- Kim, S. S., J. S. Jo, Y. J. Kim, and N. K. Sung. Authentication of rice by three-sided image analysis of kernels using two mirrors, 212
- Kim, Y. J. *See* S. S. Kim, 212
- King, G. E. *See* C. F. Morris, 16, 147
- Kiribuchi-Otobe, C., T. Nagamine, T. Yanagisawa, M. Ohnishi, and I. Yamaguchi. Production of hexaploid wheats with waxy endosperm character, 72
- Kitahara, K., T. Tanaka, T. Suganuma, and T. Nagahama. Release of bound lipids in cereal starches upon hydrolysis by glucoamylase, 1
- Klavons, J. A., F. R. Dintzis, and M. M. Millard. Hydrodynamic chromatography of waxy maize starch, 832
- *See* M. M. Millard, 687
- Klopfenstein, C. F. *See* J. E. Barrett, 168
- *See* D. K. Hix, 281
- Knuckles, B. E., W. H. Yokoyama, and M. M. Chiu. Molecular characterization of barley β -glucans by size-exclusion chromatography with multiple-angle laser light scattering and other detectors, 599
- *See* W. H. Yokoyama, 293
- Knutson, C. A. Note: Use of phenol-sulfuric acid assay to determine composition of starch-oil composites, 471
- *See* M. A. Dombink-Kurtzman, 776
- Köhler, P., B. Keck-Gassenmeier, H. Wieser, and D. D. Karsarda. Molecular modeling of the N-terminal regions of high molecular weight glutenin subunits 7 and 5 in relation to intramolecular disulfide bond formation, 154
- Kohlwey, D. *See* E. T. Champagne, 566
- Kojima, M., M. A. Hanna, and A. Gennadios. Note: Water solubility and macromolecular properties of corn meal extrudates as affected by epichlorohydrin, 526
- Köksel, H. *See* D. Dadayli, 375
- Kollengode, A. N. R., and M. A. Hanna. Flavor retention in pregelatinized and internally flavored starch extrudates, 396
- Komatsu, T. *See* R. Haginoya, 745
- Kongseree, N. *See* A. Noomhorm, 12
- Kotwal, Z. *See* B. G. Osborne, 467
- Krieger, K. M., S. A. Duvick, L. M. Pollak, and P. J. White. Note: Thermal properties of corn starch extracted with different blending methods: Microblender and homogenizer, 553
- Kruger, J. E. *See* D. W. Hatcher, 337, 502
- Kune, L. A., A. Gennadios, S. L. Cuppett, M. A. Hanna, and C. L. Weller. Cast films from soy protein isolates and fractions, 115
- Kurant, M. *See* L. C. Dickey, 676
- Kurata, T. *See* M. Nakamura, 647, 651
- Laakso, S. *See* A. Kaukovirta-Norja, 733
- Lai, H.-M., and G. W. Padua. Properties and microstructure of plasticized zein films, 771
- , —, and L. S. Wei. Properties and microstructure of zein sheets plasticized with palmitic and stearic acids, 83
- Landry, J. Note: Comparison of extraction methods for evaluating zein content of maize grain, 188
- Lang, C. E. *See* J. L. Hazelton, 400
- Larrea, M. A., M. V. E. Grossmann, A. P. Belúa, and D. Q. Tavares. Changes in water absorption and swollen volume in extruded alkaline peroxide pretreated rice hulls, 98
- Larsen, N. G. *See* W. Bushuk, 389
- Laurikainen, T. *See* J. Räsänen, 56
- Lavelli, V. *See* N. Guerrieri, 846
- Lawton, J. W., Jr. *See* W. J. Wolf, 530
- Lee, C. J., R. D. Horsley, F. A. Manthey, and P. B. Schwarz. Comparisons of β -glucan content of barley and oat, 571
- Leinen, S. D. *See* G. S. Ranhotra, 297, 361
- Leipold, H. W. *See* J. E. Barrett, 168
- Lelley, T. *See* U. Pechanek, 800
- Le Meste, M. *See* A. Rolee, 581
- Li, C.-F. *See* M.-L. Tsai, 750
- Li, W., R. Lin, and H. Corke. Physicochemical properties of common and tartary buckwheat starch, 79
- Li, W.-S., and J.-T. Shaw. Determining the fat acidity of rough rice by near-infrared reflectance spectroscopy, 556
- Lii, C.-Y. *See* S. Lu, 34
- *See* M.-L. Tsai, 750
- Lin, P.-Y., and Z. Czuchajowska. Starch properties and stability of club and soft white winter wheats from the Pacific Northwest of the United States, 639
- Lin, R. *See* W. Li, 79
- Lindborg, K. M., C. Trägårdh, A.-C. Eliasson, and P. Dejmeek. Time-resolved shear viscosity of wheat flour doughs—Effect of mixing, shear rate, and resting on the viscosity of doughs of different flours, 49
- Lingnert, H. *See* A. Rutgersson, 727
- Linscombe, S. *See* E. T. Champagne, 566
- *See* W. R. Windham, 626
- Liu, H., S. D. Arntfield, R. A. Holley, and D. B. Aime. Amylose-lipid complex formation in acetylated pea starch-lipid systems, 159
- Livesey, G. *See* R. L. Botham, 29
- Lookhart, G. L. *See* S. R. Bean, 758
- Lopes-Filho, J. F., P. Buriak, M. E. Tumbleson, and S. R. Eckhoff. Inter-mittent milling and dynamic steeping process for corn starch recovery, 633
- Lorenz, K. *See* A. Winata, 284
- Lu, S., L.-N. Chen, and C.-Y. Lii. Correlations between the fine structure, physicochemical properties, and retrogradation of amylopectins from Taiwan rice varieties, 34
- Lujan-Acosta, J., and R. G. Moreira. Effects of different drying processes on oil absorption and microstructure of tortilla chips, 216
- Lukow, O. M. *See* A. Hussain, 242, 791
- Lyon, B. G. *See* E. T. Champagne, 566
- *See* W. R. Windham, 626
- Mabesa, L. B. *See* L. S. Collado, 681
- Magnus, E. M. *See* T. Wicklund, 326
- Manthey, F. A. *See* C. J. Lee, 571
- Marchylo, B. A. *See* J. E. Dexter, 519
- Marks, B. P. *See* A. A. Perdon, 864
- *See* H. Chen, 826
- Martin, T. J. *See* W. J. Park, 7
- Masi, P. *See* L. Piazzini, 135
- Matsukawa, R. *See* R. Haginoya, 745
- Matsumoto, H. *See* M. Seguchi, 129, 384
- Matsunaga, N., and P. A. Seib. Extraction of wheat starch with aqueous sodium hydroxide, 851
- McClung, A. M. *See* E. T. Champagne, 566
- *See* W. R. Windham, 626
- McCluskey, P. J. *See* J. L. Hazelton, 400
- McDonough, C. M., B. J. Anderson, and L. W. Rooney. Structural characteristics of steam-flaked sorghum, 542
- McKenzie, K. S. *See* W. R. Windham, 626
- McKenzie, R. I. H. *See* A. Hussain, 242
- McKinstry, J. *See* V. Singh, 672
- McPherson, A. E., and P. A. Seib. Preparation and properties of wheat and corn starch maltodextrins with a low dextrose equivalent, 424
- Mehra, S. K., and S. R. Eckhoff. Single-stage short-duration tempering of corn for dry-milling, 484
- Meuser, F., N. Pahne, and M. Möller. Yield of starch and by-products in the processing of different varieties of wrinkled peas on a pilot scale, 364
- Miles, M. J. *See* P. R. Shewry, 193
- Millard, M. M., F. R. Dintzis, J. L. Willett, and J. A. Klavons. Light-scattering molecular weights and intrinsic viscosities of processed waxy maize starches in 90% dimethyl sulfoxide and H₂O, 687
- *See* J. A. Klavons, 832
- Miller, R. A., and R. C. Hoseney. Factors in hard wheat flour responsible for reduced cookie spread, 330
- , and —. Note: Use of elongational viscosity to estimate cookie diameter, 614
- , —, and C. F. Morris. Effect of formula water content on the spread of sugar-snap cookies, 669
- *See* J. L. Hazelton, 400
- Moldenhauer, K. *See* E. T. Champagne, 566
- *See* W. R. Windham, 626
- Molina-Cano, J.-L. *See* J. S. Swanston, 456
- Möller, M. *See* F. Meuser, 364
- Moore, S. A. *See* I. L. Batey, 497
- Moore, W. R. *See* D. C. Doehlert, 403
- *See* D. Zhang, 722
- Morales-Valdes, M. *See* R. I. Sánchez-Mariñez, 481
- Moreira, R. G. *See* J. Lujan-Acosta, 216
- Morgenstern, M. P. *See* A. J. Wilson, 715
- Morris, C. F., G. E. King, and G. L. Rubenthaler. Contribution of wheat flour fractions to peak hot paste viscosity, 147
- , B. J. Shackley, G. E. King, and K. K. Kidwell. Genotypic and environmental variation for flour swelling volume in wheat, 16
- *See* R. A. Miller, 669
- *See* M. Zeng, 63
- Morris, V. J. *See* R. L. Botham, 29
- Mulvaney, S. J. *See* J. Fu, 304
- Myers, D. J. *See* S. Wu, 258, 264, 268
- *See* M. Zhang, 594
- Nagahama, T. *See* K. Kitahara, 1

- Nagamine, T. See C. Kiribuchi-Otobe, 72
- Nagao, S. See R. Haginoya, 745
- Nair, R. B. See T. Demeke, 442
- Nakamura, M., and T. Kurata. Effect of L-ascorbic acid on the rheological properties of wheat flour-water dough, 647
- , and —. Effect of L-ascorbic acid and superoxide anion radical on the rheological properties of wheat flour-water dough, 651
- Nakamura, T. See T. Demeke, 442
- See K. Hayakawa, 576
- Nelsen, T. C. See F. R. Huebner, 123
- Ng, K.-Y., S. A. Duvick, and P. J. White. Thermal properties of starch from selected maize (*Zea mays* L.) mutants during development, 288
- , L. M. Pollak, S. A. Duvick, and P. J. White. Thermal properties of starch from sixty-two exotic maize (*Zea mays* L.) lines grown in two locations, 837
- Ng, P. K. W. See D. T. Campos, 489
- See R. Redaelli, 102
- Nicolas, J. See N. Kaid, 605
- Niu, Y. X. See V. Singh, 249
- Noel, T. R. See R. L. Botham, 29
- Noomhorm, A., N. Kongseree, N., and M. Apintanapong. Effect of aging on the quality of glutinous rice crackers, 12
- Obanni, M., and J. N. BeMiller. Properties of some starch blends, 431
- See M. R. Jacobson, 511
- O'Brien, L. See B. G. Osborne, 467
- Ohnishi, M. See C. Kiribuchi-Otobe, 72
- Olkku, J. See A. Kaukovirta-Norja, 733
- Ortega-Dorame, F. See R. I. Sánchez-Mariñez, 481
- Osborne, B. G., Z. Kotwal, A. B. Blakeney, L. O'Brien, S. Shah, and T. Feam. Application of the single-kernel characterization system to wheat receiving testing and quality prediction, 467
- Otto, T., B.-K. Baik, and Z. Czuchajowska. Microstructure of seeds, flours, and starches of legumes, 445
- , —, and —. Wet fractionation of garbanzo bean and pea flours, 141
- Padua, G. W. See H.-M. Lai, 83, 771
- Pahne, N. See F. Meuser, 364
- Park, H., P. A. Seib, and O. K. Chung. Fortifying bread with a mixture of wheat fiber and psyllium husk fiber plus three antioxidants, 207
- , —, —, and L. M. Seitz. Fortifying bread with each of three antioxidants, 202
- Park, W. J., D. R. Shelton, C. J. Peterson, T. J. Martin, S. D. Kachman, and R. L. Wehling. Variation in polyphenol oxidase activity and quality characteristics among hard white wheat and hard red winter wheat samples, 7
- Parris, N., L. Dickey, and J. Craig. Quantitative analysis of corn zeins by capillary electrophoresis, 766
- See L. C. Dickey, 676
- Paulsen, M. R. See K. D. Rausch, 274
- Pechanek, U., A. Karger, S. Gröger, B. Charvat, G. Schöggel, and T. Lelley. Effect of N-fertilization on quantity of flour protein components, dough properties, and breadmaking quality of wheat, 800
- Peñuelas, J. See M. Estiarte, 495
- Perdon, A. A., B. P. Marks, T. J. Siebenmorgen, and N. B. Reid. Effects of rough rice storage conditions on the amylograph and cooking properties of medium grain (cultivar Bengal) rice, 864
- Perez-Vendrell, A. See J. S. Swanson, 456
- Peri, C. See G. Giovannelli, 710
- Peterson, C. J. See W. J. Park, 7
- Piazza, L., and P. Masi. Development of crispness in cookies during baking in an industrial oven, 135
- Pogna, N. E. See R. Redaelli, 102
- Pollak, L. M., and P. J. White. Thermal starch properties in Corn Belt and exotic corn inbred lines and their crosses, 412
- See K. M. Krieger, 553
- See K.-Y. Ng, 837
- See S. K. Singh, 40
- Potus, J. See N. Kaid, 605
- Quail, K. J. See A. S. Ross, 814
- Qualset, C. O. See M. Bhattacharya, 417
- Radewonuk, E. R. See L. C. Dickey, 676
- Rakotozafy, L. See N. Kaid, 605
- Ranhotra, G. S., J. A. Gelroth, and S. D. Leinen. Effect of tristearin and other functional fats on blood lipids in hamsters, 297
- , —, —, and F. E. Schneller. Bioavailability of calcium in breads fortified with different calcium sources, 361
- Räsänen, J., J. M. V. Blanshard, M. Siitari-Kauppi, and K. Autio. Water distribution in frozen lean wheat doughs, 806
- , T. Laurikainen, and K. Autio. Fermentation stability and pore size distribution of frozen prefermented lean wheat doughs, 56
- Rausch, K. D., S. R. Eckhoff, and M. R. Paulsen. Evaluation of the displacement value as a method to detect reduced corn wet-milling quality, 274
- Redaelli, R., N. E. Pogna, and P. K. W. Ng. Effects of prolamins encoded by chromosomes 1B and 1D on the rheological properties of dough in near-isogenic lines of bread wheat, 102
- Reid, N. B. See A. A. Perdon, 864
- Reinikainen, P. See A. Kaukovirta-Norja, 733
- Reitmeier, C. A. See M. Zhang, 594
- Ring, S. G. See R. L. Botham, 29
- Robutti, J. L., F. S. Borras, and G. H. Eyherabide. Zein compositions of mechanically separated coarse and fine portions of maize kernels, 75
- Rolee, A., and M. Le Meste. Thermomechanical behavior of concentrated starch-water preparations, 581
- Rooney, L. W. See C. M. McDonough, 542
- Ross, A. S., K. J. Quail, and G. B. Crosbie. Physicochemical properties of Australian flours influencing the texture of yellow alkaline noodles, 814
- Rossnagel, B. G. See R. S. Bhatti, 190
- Rubenthaler, G. L. See C. F. Morris, 147
- Rutgersson, A., E.-L. Bergman, H. Lingnert, and A.-S. Sandberg. Optimization of temperature, time, and lactic acid concentration to inactivate lipoxygenase and lipase and preserve phytase activity in barley (cv. Blenheim) during soaking, 727
- Sakai, K. See R. Haginoya, 745
- Sánchez-Mariñez, R. I., M. O. Cortez-Rocha, F. Ortega-Dorame, M. Morales-Valdes, and M. I. Silveira. End-use quality of flour from *Rhyzopertha dominica* infested wheat, 481
- Sandberg, A.-S. See A. Rutgersson, 727
- Sapirstein, H. D. See M. G. Scanlon, 612
- Sara, R. G. See W. Bushuk, 389
- Sayre, R. N. See W. H. Yokoyama, 293
- Scanlon, M. G., D. Fahloul, and H. D. Sapirstein. Communication to the editor: A measure of fracture toughness of bread crumb, 612
- Schieberle, P. See W. Grosch, 91
- Schneeman, B. O. See W. H. Yokoyama, 293
- Schneller, F. E. See G. S. Ranhotra, 361
- Schöggel, G. See U. Pechanek, 800
- Schwarz, P. B. See C. J. Lee, 571
- Seguchi, M., M. Hayashi, and H. Matsumoto. Effect of gaseous acetic acid on dough rheological and breadmaking properties, 129
- , —, and —. Effect of a wheat protein on pH-dependent water-binding capacity and viscosity of wheat tailings fractions, 384
- , and K. Kanenaga. Study of three-dimensional structure of wheat starch granules stained with Remazolbrilliant blue dye and extracted with aqueous sodium dodecyl sulfate and mercaptoethanol solution, 548
- Seib, P. A. See N. Matsunaga, 851
- See A. E. McPherson, 424
- See H. Park, 202, 207
- See K. R. Vadlamani, 318
- Seitz, L. M. See H. Park, 202
- Shackley, B. J. See C. F. Morris, 16
- Shah, S. See B. G. Osborne, 467
- Shaw, J.-T. See W.-S. Li, 556
- Shelton, D. R. See W. J. Park, 7
- Shewry, P. R., M. J. Miles, N. H. Thomson, and A. S. Tatham. Review: Scanning probe microscopes—Applications in cereal science, 193
- Shih, F. F., and K. Daigle. Use of enzymes for the separation of protein from rice flour, 437
- Shipman, M. A. See J. L. Hazelton, 400
- Siebenmorgen, T. J. See H. Chen, 821, 826
- See A. A. Perdon, 864
- Siitari-Kauppi, M. See J. Räsänen, 806
- Silveira, M. I. See R. I. Sánchez-Mariñez, 481
- Simmons, L. D. See W. Bushuk, 389
- Singh, N. See S. K. Singh, 254
- Singh, S. K., L. A. Johnson, L. M. Pollak, S. R. Fox, and T. B. Bailey. Comparison of laboratory and pilot-plant corn wet-milling procedures, 40
- , N. Singh, L. Du, and S. R. Eckhoff. Effect of sodium hydroxide, calcium hydroxide, and potassium hydroxide on debranning of corn, 254
- See S. Wu, 264
- Singh, V., N. L. Barreiro, J. McKinstry, P. Buriak, and S. R. Eckhoff. Effect of kernel size, location, and type of damage on popping characteristics of popcorn, 672

- , and S. R. Eckhoff. Economics of germ prepreparation for dry-grind ethanol facilities, 462
- , A. E. Haken, Y. X. Niu, S. H. Zou, and S. R. Eckhoff. Hybrid-dependent effect of lactic acid on corn starch yields, 249
- Sosulski, F. W. See S. Wang, 621
- Sosulski, K. See S. Wang, 621
- Steffe, J. F. See D. T. Campos, 489
- Stephen, A. M., W. J. Dahl, D. M. Johns, and H. N. Englyst. Effect of oat hull fiber on human colonic function and serum lipids, 379
- Suárez, C. See M. P. Tolaba, 842
- Suganuma, T. See K. Kitahara, 1
- Sung, N. K. See S. S. Kim, 212
- Sünnetçioğlu, M. M. See D. Dadayli, 375
- Sutton, K. H. See W. Bushuk, 389
- Svendsen, L. See F. F. Yamin, 407
- Swanston, J. S. Waxy starch barley genotypes with reduced β -glucan contents, 452
- , R. P. Ellis, A. Perez-Vendrell, J. Voltas, and J.-L. Molina-Cano. Patterns of barley grain development in Spain and Scotland and their implications for malting quality, 456
- Takeuchi, O. See R. Haginoya, 745
- Tanaka, K. See K. Hayakawa, 576
- Tanaka, T. See K. Kitahara, 1
- Tao, H. P. See N. Zhang, 119
- Tatham, A. S. See P. R. Shewry, 193
- Tavares, D. Q. See M. A. Larrea, 98
- Thomas, K. C. See S. Wang, 621
- Thompson, D. B. See D. K. Fisher, 344
- Thomson, N. H. See P. R. Shewry, 193
- Tilley, K. A. Detection of *O*-glycosidically linked mannose within the structure of a highly purified glutenin subunit isolated from Chinese spring wheat, 371
- Tolaba, M. P., R. J. Aguerre, and C. Suárez. Modeling cereal grain drying with variable diffusivity, 842
- Trägårdh, C. See K. M. Lindborg, 49
- Tsai, M.-L., C.-F. Li, and C.-Y. Lii. Effects of granular structures on the pasting behaviors of starches, 750
- Tumbleson, M. E. See J. F. Lopes-Filho, 633
- Turnlund, J. R. See W. H. Yokoyama, 293
- Tyler, R. T. See T. Vasanthan, 25
- Vadlamani, K. R., and P. A. Seib. Two metal ions improve brightness in wheat-dough products and affect aqueous dispersion of gluten, 318
- Vasanthan, T., R. S. Bhatta, R. T. Tyler, and P. Chang. Isolation and cationization of barley starches at laboratory and pilot scale, 25
- Vensel, W. H., A. E. Adalsteins, and D. D. Kasarda. Purification and characteristics of the glutenin subunits of *Triticum tauschii*, progenitor of the D genome in hexaploid bread wheat, 108
- Villareal, C. P., S. Hizukuri, and B. O. Juliano. Amylopectin staling of cooked milled rices and properties of amylopectin and amylose, 163
- Vinyard, B. T. See E. T. Champagne, 566
- Voltas, J. See J. S. Swanston, 456
- Walker, C. E. See J. L. Hazelton, 400
- , See D. K. Hix, 281
- Wang, S., K. C. Thomas, W. M. Ingledew, K. Sosulski, and F. W. Sosulski. Rye and triticale as feedstock for fuel ethanol production, 621
- Watts, B. M. See A. Hussain, 242
- Webb, B. D. See E. T. Champagne, 566
- , See W. R. Windham, 626
- Wehling, R. L. See W. J. Park, 7
- Wehrle, K., H. Grau, and E. K. Arendt. Effects of lactic acid, acetic acid, and table salt on fundamental rheological properties (mixing time and rest time) of wheat dough, 739
- Wei, L. S. See H.-M. Lai, 83
- Weisz, J. See M. U. Beer, 476, 705
- Weller, C. L. See R. A. Buffo, 473
- , See L. A. Kunte, 115
- White, P. J. See K. M. Krieger, 553
- , See K.-Y. Ng, 288, 837
- , See L. M. Pollak, 412
- , See F. F. Yamin, 407
- Wicklund, T., and E. M. Magnus. Effect of extrusion cooking on extractable lipids and fatty acid composition in sifted oat flour, 326
- Wieser, H. See P. Köhler, 154
- Willet, J. L. See M. M. Millard, 687
- Wilson, A. J., A. R. Wooding, and M. P. Morgenstern. Comparison of work input requirement on laboratory-scale and industrial-scale mechanical dough development mixers, 715
- Wilson, J. D. See D. B. Bechtel, 235
- Winata, A., and K. Lorenz. Effects of fermentation and baking of whole wheat and whole rye sourdough breads on cereal alkylresorcinols, 284
- Windham, W. R., B. G. Lyon, E. T. Champagne, F. E. Barton, II, B. D. Webb, A. M. McClung, K. A. Moldenhauer, S. Linscombe, and K. S. McKenzie. Prediction of cooked rice texture quality using near-infrared reflectance analysis of whole-grain milled samples, 626
- Wolf, W. J., and J. W. Lawton, Jr. Isolation and characterization of zein from corn distillers' grains and related fractions, 530
- Wood, P. J. See M. U. Beer, 476, 705
- Wooding, A. R. See A. J. Wilson, 715
- Wrigley, C. W. See M. Zeng, 63
- Wu, S., D. J. Myers, and L. A. Johnson. Effects of maize hybrid and meal drying conditions on yield and quality of extracted zein, 268
- , —, and —. Factors affecting yield and composition of zein extracted from commercial corn gluten meal, 258
- , —, —, S. R. Fox, and S. K. Singh. Note: Pilot-plant wet-milling process for producing corn gluten meal, 264
- Yamaguchi, I. See C. Kiribuchi-Otobe, 72
- Yamin, F. F., L. Svendsen, and P. J. White. Thermal properties of corn starch extraction intermediates by differential scanning calorimetry, 407
- Yanagisawa, T. See C. Kiribuchi-Otobe, 72
- Yokoyama, K. See R. Haginoya, 745
- Yokoyama, W. H., C. A. Hudson, B. E. Knuckles, M.-C. M. Chiu, R. N. Sayre, J. R. Turnlund, and B. O. Schneeman. Effect of barley β -glucan in durum wheat pasta on human glycemic response, 293
- , See B. E. Knuckles, 599
- Zeng, M., C. F. Morris, I. L. Batey, and C. W. Wrigley. Sources of variation for starch gelatinization, pasting, and gelation properties in wheat, 63
- Zhang, D., D. C. Doehlert, and W. R. Moore. Factors affecting viscosity of slurries of oat groat flours, 722
- Zhang, M., C. A. Reitmeier, E. G. Hammond, and D. J. Myers. Production of textile fibers from zein and a soy protein-zein blend, 594
- Zhang, N., B. L. Jones, and H. P. Tao. Purification and characterization of a new class of insect α -amylase inhibitors from barley, 119
- Zou, S. H. See V. Singh, 249

Subject Index

Page numbers of errata are in italics.

Absorption, prediction; regression equation for (Hazelton et al), 400

Alkaline

—hydrogen peroxide extraction; for hemicellulose isolation from corn fiber (Doner and Hicks), 176

—processing; to detoxify rapeseed meal (Barrett et al), 168

Amino acid, sequencing; of *Triticum tauschii* (Vensel et al), 108

α -Amylase, inhibitors; in barley (Zhang et al), 119

Amylopectin

—hydrodynamic chromatography of waxy maize starch for molecular weight values (Klavons et al), 832

—molecular weight and intrinsic viscosity (Millard et al), 687

—retrogradation; correlation with fine structure and physicochemical properties from Taiwan rice (Lu et al), 34

—staling of cooked milled rice (Villareal et al), 163

Amylose, zero lines, hull-less barley (Bhatty and Rossnagel), 190

Antioxidants

—bread fortification with fiber and (Park et al), 207

—for fortifying bread (Park et al), 202

—naturally occurring; cookie physical and chemical attributes and consumer acceptance (Hix et al), 281

L-Ascorbic acid, effect on rheological properties of wheat flour-water dough (Nakamura and Kurata), 647, 651

Baking

—cookies: effect of formula water content on spread (Miller et al), 669; hard wheat factors responsible for reduced spread (Miller and Hosney), 330

—effects of whole wheat and rye sourdough breads on cereal alkylresorcinols (Winata and Lorenz), 284

—elongational viscosity to estimate cookie diameter (Miller and Hosney), 614

—quality: extra strong wheat (communication to the editor) (Scanlon et al), 612; influenced by kernel size and shriveling (Gaines et al), 700

—temperature effect on crumb staling (Giovannelli et al), 710

Barley

—carbohydrates resistant to digestion in human ileostomate; physicochemical characterization (Botham et al), 29

—compared to barley, β -glucan content (Lee et al), 571

—consecutive extracts of, molecular weight distribution and β -glucan content of (Beer et al), 476

— β -glucan, effect on glycemic response (Yokoyama et al), 293

—grain development patterns; implications for malting quality (Swanston et al), 456

—hull-less: milling for bran and flour production (Bhatty), 693; zero amylose lines (Bhatty and Rossnagel), 190

—insect α -amylase inhibitors (Zhang et al), 119

—lipoxygenase and lipase inactivation and phytase preservation in (Rutgers et al), 727

—molecular characterization of β -glucans by size exclusion chromatography with laser light scattering detection (Knuckles et al), 599

—starch lipid modification by malting (Kaukovirta-Norja et al), 733

—starches; isolation and cationization of (Vasanthan et al), 25

—waxy starch genotypes (Swanston), 452

Bioavailability, of calcium; in breads fortified with different calcium sources (Ranhotra et al), 361

Bran

—production; hull-less barley milling for (Bhatty), 693

—rice, protein fraction characterization of (Hamada), 662

Bread

—baking temperature effect on crumb-staling kinetics (Giovannelli et al), 710

—calcium bioavailability; fortification with different calcium sources (Ranhotra et al), 361

—crumb strength (communication to the editor) (Scanlon et al), 612

—fiber and antioxidants for fortification (Park et al), 207

—fortification with antioxidants (Park et al), 202

— γ -gliadin related to loaf volume (Huebner et al), 123

—whole wheat and rye sourdough, effects of fermentation and baking on cereal alkylresorcinols (Winata and Lorenz), 284

Breadmaking

—properties; affected by gaseous acetic acid (Seguchi et al), 129

—quality: individual subunit effect on (Huang and Khan), 786; quantitative effect of total amounts on (Huang and Khan), 781; of wheat affected by N-fertilization (Pechane'x et al), 800

Buckwheat, common and tartary, physicochemical properties (Li et al), 78

Calcium, bioavailability; in breads fortified with different calcium sources (Ranhotra et al), 361

Carbohydrates, liquifaction of, for protein separation from rice flour (Shih and Daigle), 437

Cereal grain

—drying (Tolaba et al), 842

—structure and composition (Irving and Jideani), 224

Chromatography, hydrodynamic, amylopectin, for molecular weight determination (Klavons et al), 832

Color, measurement of noodles; automated system for (Corke et al), 356

Cookies

—containing naturally occurring antioxidants, physical and chemical attributes and consumer acceptance (Hix et al), 281

—crispness development during industrial oven baking (Piazza and Masi), 135

—effect of formula water content on spread (Miller et al), 669

—hard wheat factors responsible for reduced spread (Miller and Hosney), 330

Corn (see also Maize)

—debranning; effect of sodium, calcium, and potassium hydroxides (Singh et al), 254

—dry-milled: *Fusarium* mold and fumonisin distribution in (Katta et al), 858; hydrocyclone separation of (Dickey et al), 676

—extraction of zein (Wolf and Lawton), 530

—fiber: hemicellulose isolation by alkaline hydrogen peroxide extraction (Doner and Hicks), 176; recovery of starch and protein (Dowd), 589

—flour and starch; lime effect on gelatinization of (Bryant and Hamaker), 171

—gluten meal; effect of drying (Wu et al), 258

—single-stage short-duration tempering of, for dry milling (Mehra and Eckhoff), 484

—starch granules; visualization of channels and cavities of (Huber and BeMiller), 537

—wet milling, effect of processing conditions (Wu et al), 264, 268

—wet-milling procedures; comparison of laboratory and pilot-plant (Singh et al), 40

—wet-milling quality and prediction (Rausch et al), 274

—zeins; capillary electrophoresis for quantitative analysis (Parris et al), 766

Corn meal

—extrudates affected by epichlorohydrin (Kojima et al), 526

—particle size (Garber et al), 656

Corn starch

—DSC, thermal analysis, starch extraction (Yamin et al), 407

—extraction and blending methods (Krieger et al), 553

—hybrid-dependent effect of lactic acid on yield (Singh et al), 249

—malodextrins with low dextrose equivalent; preparation and properties (McPherson and Seib), 424

—recovery by milling and steeping (Lopes-Filho et al), 633

—thermal properties; changes during growth development (Ng et al), 288

Debranning, of corn; effect of sodium, calcium, and potassium hydroxides (Singh et al), 254

Detoxification, of rapeseed meal by alkaline extrusion (Barrett et al), 168

Dietary fiber

—effect on glycemic response (Yokoyama et al), 293

—effect on human colonic function and serum lipids (Stephen et al), 379

Differential scanning calorimetry, corn starch, thermal analysis, starch extraction (Yamin et al), 407

Dough

—frozen lean wheat, water distribution in (Räsänen et al), 806

—mechanical, effect on extractability of wheat storage proteins from bread dough (Bushuk et al), 389

—mixing requirement on laboratory and industrial scale (Wilson et al), 715

—mixing strength (Hussain et al), 242

—prefermented, fermentation stability and pore structure (Räsänen et al), 56

—rheology; affected by gaseous acetic acid (Seguchi et al), 129

—wheat: brightness affected by metal ions (Vadlamani and Seib), 318; effects of lactic acid, acetic acid, and salt on rheological properties of (Wehrle et al), 739

—wheat flour: rheological properties (Fu et al), 304; time-resolved shear viscosity of (Lindborg et al), 49

Dry milling

—composition of oat bran and flour prepared by (Doehlert and Moore), 403

—corn for, single-stage short-duration tempering of (Mehra and Eckhoff), 484

Drying

- cereal grain (Tolaba et al), 842
- of corn gluten meal (Wu et al), 264, 268
- effect on oil absorption and microstructure of tortilla chips (Lujan-Acosta and Moreira), 216

Economics, of germ prepreparation; for dry-grind ethanol facilities (Singh and Eckhoff), 462

Electrophoresis

- capillary combined with HPLC 2-D separation (Bean and Lookhart), 758
- capillary, for quantitative analysis of corn zeins (Parris et al), 766
- multistacking SDS-PAGE of glutenin (Huang and Khan), 229

Endosperm, structure of wheat starchy endosperm dried by different methods (Bechtel and Wilson), 235

Enzymes, α -amylase, cellulase, and hemicellulase, in processing of rice flour for protein separation (Shih and Daigle), 437

Errata

- vol. 73, no. 5, p. 574, 90
- vol. 73, no. 6, p. 777, 359

Ethanol

- dry-grind facilities, economics of germ prepreparation (Singh and Eckhoff), 462

—fuel production; rye and triticale as feedstock for (Wang et al), 621

Extractability, of wheat storage proteins from bread dough; effect of mechanical dough development on (Bushuk et al), 389

Extrusion

- alkaline, to detoxify rapeseed meal (Barrett et al), 168
- changes in water absorption and swollen volume in alkaline peroxide pretreated rice hulls by (Larrea et al), 98
- corn meal, epichlorohydrin effect on water solubility and macromolecular properties (Kojima et al), 526

Extrusion cooking

- effect of corn meal particle size (Garber et al), 656
- effect on extractable lipids and fatty acid composition in oat flour (Wicklund and Magnus), 326

Fat acidity, of rough rice; determination by NIR spectroscopy (Li and Shaw), 556

Fermentation, effects of whole wheat and rye sourdough breads on cereal alkylresorcinols (Winata and Lorenz), 284

Fiber

- bread fortification with antioxidants and (Park et al), 207
- textile, production from zein and soy protein-zein blend (Zhang et al), 594

Film

- cast, from soy protein isolates and fractions (Kunte et al), 115
- physical properties for those produced using zein and kafirin (Buffo et al), 473

Flavor

- of cereal products, review (Grosch and Schieberle), 91
- retention in starch extrudates (Kollengode and Hanna), 396

Flour

- Australian, influencing texture of yellow alkaline noodles (Ross et al), 814
- composition; dry-milling mechanisms of (Doehler and Moore), 403
- end-use quality of; from *Rhyzopertha dominica* infested wheat (Sánchez-Mariñez et al), 481
- hard wheat, reduced cookie spread factors (Miller and Hosney), 330
- legume, wet fractionation (Otto et al), 141
- oat, extrusion cooking effect on extractable lipids and fatty acid composition (Wicklund and Magnus), 326
- production; hull-less barley milling for (Bhatty), 693
- simple phenolic acids in, relation to ash content, color, and polyphenol oxidase activity (Hatcher and Kruger), 337
- swelling volume in wheat; genotypic and environmental variation (Morris et al), 16
- wheat: contribution to peak hot paste viscosity (Morris et al), 147; glutathione-dehydroascorbate oxidoreductase studies (Kaid et al), 605; mixing characteristics influenced by gliadin-rich subfractions of Glenlea wheat (Hussain and Lukow), 791; relation between laboratory and industrial scale mixing (Wilson et al), 715
- yield; predicting with single-kernel characterization system (Osborne et al), 467

Fractionation, of glutenin by multistacking SDS-PAGE (Huang and Khan), 229

Gelatinization

- of corn flour and starch; lime effect (Bryant and Hamaker), 171
- starch, in wheat (Zeng et al), 63

Gliadin

- γ -gliadin correlated to loaf volume (Huebner et al), 123
- of near-isogenic bread lines, effects on dough properties (Redaelli et al), 102

β -Glucan

- in barley, effect on glycemic response (Yokoyama et al), 293
- comparison of barley and oat (Lee et al), 571
- content of consecutive extracts of oat and barley cultivars, molecular weight distribution (Beer et al), 476
- molecular characterization by size exclusion chromatography with laser light scattering detection (Knuckles et al), 599
- from oats, cooking and storage effect on (Beer et al), 705
- reduced, in waxy starch barley genotypes (Swanston), 452

Glucosylase, hydrolysis; bound lipid release in cereal starches (Kitahara et al), 1

Glutathione, dehydroascorbate oxidoreductase from wheat flour (Kaid et al), 605

Gluten

- aqueous dispersion affected by metal ions (Vadlamani and Seib), 318
- oxidant action effects (Cherian and Chinachoti), 312

Glutenin

- characterization by multistacking SDS-PAGE (Huang and Khan), 229
- HMW subunits; molecular modeling (Köhler et al), 154
- of near-isogenic bread lines, effects on dough properties (Redaelli et al), 102
- subunit isolated from Chinese spring wheat; detection of *O*-glycosidically linked mannose (Tilley), 371
- subunits of *Triticum tauschii*, purification and characterization (Vensel et al), 108

Grain

- color classification of (Dowell), 617
- maize, extraction methods for evaluating zein content (Landry), 188

Hardness

- maize endosperm, related to amylose content and damage susceptibility (Dombrink-Kurtzman and Knutson), 776
- of maize kernels as influenced by zein composition (Robutti et al), 75
- wheat endosperm structure and its relationship to endosperm texture (Bechtel and Wilson), 235
- Hemicellulose, isolation from corn fiber; by alkaline hydrogen peroxide extraction (Doner and Hicks), 176

HPLC

- combined with capillary electrophoresis 2-D separation (Bean and Lookhart), 758
- RP- and size-exclusion to quantitate protein fractions relating to breadbaking quality (Huebner et al), 123
- zein composition of coarse and fine portions of ground kernels by RP- (Robutti et al), 75

Image analysis, with three sides of rice kernel for authentication (Kim et al), 212

Insects, α -amylases inhibited by barley proteins (Zhang et al), 119

Instructions to authors, iv

Irradiation, detection in wheat, using EPR spin probe technique (Dadayli et al), 375

Kafirin, films; comparison to zein films (Buffo et al), 473

Lactic acid, on corn starch yields; hybrid-dependent effect (Singh et al), 249

Legumes

- microstructure of (Otto et al), 445
- wet fractionation (Otto et al), 141
- Lipase, inactivation; in barley (Rutgersson et al), 727

Lipids

- amylose complexes in modified starch (Liu et al), 159
- in barley and malt starches (Kaukovirta-Norja et al), 733
- blood, effect of tristearin and other functional fats on (Ranhotra et al), 297
- extractable, effect of extrusion cooking on (Wicklund and Magnus), 326
- release of bound, in cereal starches (Kitahara et al), 1
- serum, oat hull fiber effect on (Stephen et al), 379
- Lipoxygenase, inactivation; in barley (Rutgersson et al), 727

Maize (see also Corn)

- endosperm hardness related to amylose content and damage susceptibility (Dombrink-Kurtzman and Knutson), 776
- endosperm mutants; thermal property changes during maturity stages (Ng et al), 288
- grain; extraction methods for evaluating zein content (Landry), 188
- thermal properties of starch from (Ng et al), 837
- whole grain amylose analysis, using NITS (Campbell et al), 300
- zein composition of coarse and fine portions of ground kernels (Robutti et al), 75

Malting

- effect on starch lipids (Kaukovirta-Norja et al), 733
- quality affected by barley grain development patterns (Swanston et al), 456

Maltodextrins, corn starch, preparation and properties (McPherson and Seib), 424

Methods

- blending, for corn starch extraction (Krieger et al), 553
- extraction of zein for evaluating its content (Landry), 188
- flow injection analysis biosensor, for damaged starch determination (Haginoya et al), 745

Microscopy, pore size distribution of frozen doughs (Räsänen et al), 56

Microstructure

- cereal grain, *Digitariaeaxis* (Irving and Jideani), 224
- drying effect on, in tortilla chips (Lujan-Acosta and Moreira), 216
- of garbanzo beans and peas (Otto et al), 445

Milling

- corn fiber (Dowd), 589
- for corn starch recovery (Lopes-Filho et al), 633
- dry: of corn fractions, *Fusarium* mold and fumonisin distribution (Katta et al), 858; hydrocyclone separation of corn (Dickey et al), 676
- effect on rice flavor (Champagne et al), 566
- of hull-less barleys for bran and flour production (Bhatti), 693
- rice kernel thickness effect on (Chen and Siebenmorgen), 821
- roller mill extraction rate effect on chapatti-making quality (Hatcher et al), 502

- semolina, *Fusarium* head blight effect on (Dexter et al), 519
- soft wheat, influence of kernel size and shriveling (Gaines et al), 700

Mixograph, regression equation for predicting absorption for (Hazelton et al), 400

Moisture, effect on rice flavor (Champagne et al), 566

Molecular modeling, of N-terminal regions of HMW glutenin subunits (Köhler et al), 154

Near-infrared transmittance spectroscopy, for whole grain amylose analysis in maize (Campbell et al), 300

NIR spectroscopy

- for determining fat acidity of rough rice (Li and Shaw), 556
- for prediction of cooked rice texture (Windham et al), 626
- for quantifying surface lipid content of milled rice (Chen et al), 826
- using visible wavelengths to classify wheat (Dowell), 617

Nonprotein nitrogen, of corn flour, gluten meal and distillers' grains (Wolf and Lawton), 530

Nonstarchy polysaccharides, content and composition of, in sorghum endosperms (Kavitha and Chandrashekar), 22

Noodles

- color; continuous measurement of time-dependent changes (Corke et al), 356
- genetic variation in color of sweetpotato flour (Collado et al), 681
- quality; prediction by RVA viscosity (Batey et al), 497
- starch, properties affected by sweetpotato genotype (Collado and Corke), 182
- yellow alkaline, texture influenced by Australian flours (Ross et al), 814

Oat

- compared to barley, β -glucan content (Lee et al), 571
- consecutive extracts of, molecular weight distribution and β -glucan content of (Beer et al), 476
- cooking and storage effect on molecular weight of β -glucan extracted from (Beer et al), 705

—groat flours; factors affecting viscosity of slurries (Zhang et al), 722

Oat bran, composition; dry-milling mechanisms of (Doehle and Moore), 403

Oil

- absorption; drying effect on, in tortilla chips (Lujan-Acosta and Moreira), 216
- images of oil bodies (Shewry et al), 193

Oxidation, action effects on gluten (Cherian and Chinachoti), 312

Pasta

- durum wheat, *Fusarium* head blight effect on (Dexter et al), 519
- high fiber, glycemic properties (Yokoyama et al), 293

Phenolic acids, in flours from Canadian wheat; relation to ash content, color, and polyphenol oxidase activity (Hatcher and Kruger), 337

Phenol-sulfuric acid assay, determination of composition of starch-oil composites (Knutson), 471

Physicochemical properties

- of barley carbohydrates resistant to digestion in human ileostomate (Botham et al), 29
- of common and tartary buckwheat (Li et al), 78

—correlations, fine structure and amylopectin retrogradation from Taiwan rice (Lu et al), 34

Polyphenol oxidase, variation, in wheat (Park et al), 7

Popcorn, effect of kernel size, location, and type of damage on popping characteristics (Singh et al), 672

Protein

- corn, capillary electrophoresis for zein quantitative analysis (Parris et al), 766
- corn protein extraction (Wu et al), 258, 264, 268
- images of gliadin and glutenins (Shewry et al), 193
- interaction with starch (Guerrieri et al), 846
- separation from rice flour, using carbohydrate-hydrolyzing enzymes (Shih and Daigle), 437
- wheat: effects of chromosomes 1B and 1D on dough properties (Redaelli et al), 102; separation by combined HPLC-CE (Bean and Lookhart), 758; storage from bread dough, effect of mechanical dough development on extractability (Bushuk et al), 389

Retrogradation

- of maize starch (Fisher and Thompson), 344
- resistance to, waxy hexaploid wheat starch (Hayakawa et al), 576

Review

- flavor of cereal products (Grosch and Schieberle), 91
- scanning probe microscopes, cereal science applications (Shewry et al), 193

Reviewers, acknowledgment of, iii

Rheological properties

- of concentrated starch-water preparations (Rolee and Le Meste), 581
- effects of wheat prolamins encoded by chromosomes 1B and 1D (Redaelli et al), 102
- of full-formula doughs (Hussain et al), 242
- of wheat flour doughs, added fat effect (Fu et al), 304
- of wheat flour-water dough: effect of L-ascorbic acid (Nakamura and Kurata), 647; effect of L-ascorbic acid and superoxide anion radical (Nakamura and Kurata), 651

Rheology

- cookie crispness development during industrial oven baking (Piazza and Masi), 135
- cookie dough (Miller and Hoseney), 614
- dough, affected by gaseous acetic acid (Seguchi et al), 129
- of undeveloped and developed wheat dough (Campos et al), 489

Rice

- authentication by image analysis (Kim et al), 212
- bran; protein fraction characterization of (Hamada), 662
- cooked milled, amylopectin staling of (Villareal et al), 163
- cooked texture prediction using NIR analysis (Windham et al), 626
- correlations between fine structure, physicochemical properties, and amylopectin retrogradation (Lu et al), 34
- crackers; aging effect on quality of (Noomhorm et al), 12
- flavor; effects of drying, moisture content, and milling degree on (Champagne et al), 566
- hulls; extrusion of alkaline peroxide pretreated (Larrea et al), 98
- kernel thickness effect on milling degree (Chen and Siebenmorgen), 821
- low-glutenin content mutant; properties and protein quality (Eggum and Juliano), 200
- medium grain amylograph and cooking properties affected by rough rice storage conditions (Perdon et al), 864
- milled, lipid content quantification via visible/NIR spectroscopy (Chen et al), 826
- rough, fat acidity determination by NIR spectroscopy (Li and Shaw), 556
- starch-lipid complex effect on in vitro digestibility, complexing index, and viscosity (Guraya et al), 561
- Roselle (Karkade), protein fractionation and functional properties (Abu-Tarboush et al), 352
- Rye, as feedstock for fuel ethanol production (Wang et al), 621

Scanning electron microscopy, of *Digitariaeaxis* grain (Irving and Jideani), 224

SDS-PAGE

- for quantitative determination of HMW glutenin subunits of HRS wheat (Huang and Khan), 781, 786
- of *Triticum tauschii* (Vensel et al), 108

Sorghum

- content and composition of nonstarchy polysaccharides in endosperms of (Kavitha and Chandrashekar), 22
- kafirin, physical properties of films (Buffo et al), 473
- starch granules, visualization of channels and cavities of (Huber and BeMiller), 537
- value in steam flaking (McDonough et al), 542

- Sourdough, whole wheat and rye, effects of fermentation and baking on cereal alkylresorcinols (Winata and Lorenz), 284
- Soy protein
- isolates and fractions; cast films from (Kunte et al), 115
 - textile fiber production from (Zhang et al), 594
- Staling, baking temperature effects on (Giovannelli et al), 710
- Starch
- amylose-lipid complexes in modified starch, DSC measurement (Liu et al), 159
 - barley, isolation and cationization of (Vasanthan et al), 25
 - blends; properties of (Obanni and BeMiller), 431
 - buckwheat, common and tartary (Li et al), 78
 - cereal, bound lipid release in, upon glucoamylase hydrolysis (Kitahara et al), 1
 - corn thermal properties (Pollak and White), 412
 - damage: determination by flow injection analysis biosensor method (Haginoya et al), 745; predicting with single-kernel characterization system (Osborne et al), 467
 - exudates; flavor retention in (Kollengode and Hanna), 396
 - gelatinization, pasting, and gelation properties in wheat (Zeng et al), 63
 - granules of corn and sorghum; visualization of channels and cavities of (Huber and BeMiller), 537
 - interaction with protein (Guerrieri et al), 846
 - lime effect on gelatinization of (Bryant and Hamaker), 171
 - lipids; effect of malting on (Kaukovirta-Norja et al), 733
 - maize, retrogradation (Fisher and Thompson), 344
 - molecular weight and intrinsic viscosity (Millard et al), 687
 - pasting; wheat flour fraction contribution (Morris et al), 147
 - pasting behaviors; effects of granular structures on (Tsai et al), 750
 - pasting properties; Iranian hexaploid wheat landraces (Bhattacharya et al), 417
 - properties of gelatinization and retrogradation of waxy hexaploid wheat (Hayakawa et al), 576
 - properties and stability of club and soft white winter wheats (Lin and Czuchajowska), 639
 - recovery from corn fiber (Dowd), 589
 - retrogradation of (Jacobson et al), 511
 - rice, starch-lipid complex effect on in vitro digestibility, complexing index, and viscosity (Guraya et al), 561
 - role in flake quality (McDonough et al), 542
 - starch-oil composites; determination of composition (Knutson), 471
 - sweetpotato, properties of noodles (Collado and Corke), 182
 - thermal properties; from maize (Ng et al), 837
 - viscosity; variation caused by operating conditions and effects on prediction of noodle quality (Batey et al), 497
 - water preparations; thermomechanical behavior of (Rolee and Le Meste), 581
 - waxy barley genotypes (Swanston), 452
 - wheat, extraction with aqueous sodium hydroxide (Matsunaga and Seib), 851
 - yield; purify of products (Meuser et al), 364
- Steam flaking, quality of flakes (McDonough et al), 542
- Storage, rough rice, effects on amylograph and cooking properties of medium grain rice (Perdon et al), 864
- Sweetpotato
- flour; color genetic variation related to use in wheat-based flour products (Collado et al), 681
 - genotype; starch noodle properties affected by (Collado and Corke), 182
- Techniques, scanning probe microscopes (Shewry et al), 193
- Thermal properties, maize mutants, effect of endosperm maturity on (Ng et al), 288
- Tortillas, oil absorption and microstructure of, drying effect on (Lujan-Acosta and Moreira), 216
- Tricin, separation from wheat leaves; RP-HPLC method for (Estiarte et al), 495
- Tristearin, effect on blood lipids in hamsters (Ranhotra et al), 297
- Triticale, as feedstock for fuel ethanol production (Wang et al), 621
- Viscosity
- of oat groat flours; factors (Zhang et al), 722
 - time-resolved shear, of wheat flour doughs (Lindborg et al), 49
 - wheat tailings fractions; effect of wheat protein on (Seguchi et al), 384
- Wet milling
- corn: comparison of laboratory and pilot-plant procedures (Singh et al), 40; lactic acid effect (Singh et al), 249
 - laboratory milling procedures (Rausch et al), 274
 - of wrinkled peas (Meuser et al), 364
- Wheat
- breadmaking quality affected by N-fertilization (Pechanek et al), 800
 - Chinese spring, detection of *O*-glycosidically linked mannose within glutenin subunit (Tilley), 371
 - club and soft white winter, starch properties and stability of (Lin and Czuchajowska), 639
 - dough: effects of lactic acid, acetic acid, and salt on rheological properties of (Wehrle et al), 739; fermentation stability and pore size distribution after frozen storage (Räsänen et al), 56
 - dough rheology (Fu et al), 304; (Campos et al), 489
 - durum, *Fusarium* head blight effect on semolina milling and pasta-making quality of (Dexter et al), 519
 - evaluation for waxy proteins (Demeke et al), 442
 - flour: determination of diastatic activity in by biosensor (Haginoya et al), 745; fractions; contribution to peak hot paste viscosity (Morris et al), 147; glutathione-dehydroascorbate oxidoreductase studies (Kaid et al), 605; mixing characteristics influenced by gliadin-rich subfractions of Glenlea wheat (Hussain and Lukow), 791; swelling volume in, genotypic and environmental variation (Morris et al), 16
 - glutenin of hard red spring (Huang and Khan), 229
 - grain; single-kernel characterization technology applied to (Osborne et al), 467
 - hexaploid; Iranian landraces (Bhattacharya et al), 417; production with waxy endosperm character (Kiribuchi-Otobe et al), 72
 - irradiated, detection using EPR spin probe technique (Dadayli et al), 375
 - leaves; tricin separation from, RP-HPLC method for (Estiarte et al), 495
 - NaOH test for color classification (Dowell), 614
 - protein fractions relating to bread-baking quality (Huebner et al), 123
 - quality characteristics of waxy hexaploid (Hayakawa et al), 576
 - Rhizopertha dominica* infested, end-use quality of flour from (Sánchez-Mariñez et al), 481
 - soft, milling and baking quality influenced by kernel size and shriveling (Gaines et al), 700
 - starch; extraction with aqueous sodium hydroxide (Matsunaga and Seib), 851
 - starch gelatinization, pasting, and gelation properties in (Zeng et al), 63
 - starch granules; stained with Remazolbrilliant blue dye (Seguchi and Kanenaga), 548
 - tailings; effect of wheat protein on water-binding capacity and viscosity of (Seguchi et al), 384
 - ultrastructure of developing endosperm dried by different methods (Bechtel and Wilson), 235
 - variation in polyphenol oxidase activity and quality characteristics (Park et al), 7
- Zein
- biodegradable sheets based on, plasticized with palmitic and stearic acids (Lai et al), 82
 - composition: comparison between coarse and fine portion of ground kernels (Robutti et al), 75; factors affecting (Wu et al), 258
 - extraction: from corn flour, gluten meal, and distillers' grains (Wolf and Lawton), 530; factors affecting (Wu et al), 268; methods for evaluating its content (Landry), 188
 - films; comparison to kafirin films (Buffo et al), 473
 - plasticized films; properties and microstructure of (Lai and Padua), 771
 - textile fiber production from (Zhang et al), 594